

A Patient's Guide to **Trochanteric Bursitis of the Hip**




Iain is a specialist in musculoskeletal imaging and the diagnosis of musculoskeletal pain. This information is provided with the hope that you can better understand and manage your condition. The information is not specific to your condition and is meant as a general guide only. Iain has added some information at the end of each booklet which might help add some context in regard to assessment and management from a local perspective.

DR IAIN DUNCAN

info@driainduncan.com.au

All materials within these pages are the sole property of Medical Multimedia Group, LLC and are used herein by permission. eOrthopod is a registered trademark of Medical Multimedia Group, LLC.



Introduction

A common spot for *bursitis* is on the side of the hip. Here a large tendon passes over the bony bump on the side of the hip. The bony bump is called the *greater trochanter*. Inflammation in the bursa between the tendon and the greater trochanter is called *trochanteric bursitis*. This problem is common in older individuals. It may also occur in younger patients who are extremely active in exercises such as walking, running, or biking.

This guide will help you understand


- how trochanteric bursitis develops
- how doctors diagnose the condition
- what treatments are available

Anatomy

Where is the trochanteric bursa, and what does it do?

The hip joint is one of the true ball-and-socket joints of the body. The hip socket is called the *acetabulum* and forms a deep cup that surrounds the ball of the upper thigh bone (*femur*), or *femoral head*. Thick muscles of the buttock at the back and the thick muscles of the thigh in the front surround the hip.

The greater trochanter is the large bump on the outside of the upper end of the femur. This bump is the point where the large buttock muscles that move the hip connect to the femur. The *gluteus maximus* is the largest of these muscles. It attaches lower down on the femur.




Where friction occurs between muscles, tendons, and bones, there is usually a structure called a *bursa*. A bursa is a thin sac of tissue that contains fluid to lubricate the area and reduce friction. The bursa is a normal structure. The body will even produce a bursa in response to friction.

Causes

Why do I have this problem?


Sometimes a bursa can become *inflamed* (swollen and irritated) because of too much friction or because of an injury to the bursa. An inflamed bursa can cause pain because movement makes the structures around the bursa rub against it.

Friction can build in the bursa during walking if the long tendon on the side of the thigh is tight. It is unclear what causes this tightening of the tendon. The gluteus maximus attaches to this long tendon. As you walk, the gluteus maximus pulls this tendon over the greater



trochanter with each step. When the tendon is tight, it rubs against the bursa. The rubbing causes friction to build in the bursa, leading to irritation and inflammation. Friction can also start if the outer hip muscle (*gluteus medius*) is weak, if one leg is longer than the other, or if you run on banked (slanted) surfaces.

Most cases of trochanteric bursitis appear gradually with no obvious underlying injury or cause. Trochanteric bursitis can occur after artificial replacement of the hip joint or other types of hip surgery. The cause may be a combination of changes in the way the hip works, the way it is aligned, or the way scar tissue has formed from the healing incision.




A **fall on the hip** can cause bleeding into the bursa, forming a *hematoma*. The bleeding is not serious, but the bursa may react to the blood by becoming inflamed. The inflammation causes the bursa to become thickened over time. This thickening, constant irritation, and inflammation may result in the condition becoming chronic, or long lasting.

Symptoms

What does the condition feel like?

The first symptom of trochanteric bursitis is usually pain. The pain can be felt in the area of the hip right over the bump that forms the greater trochanter. Eventually the pain may radiate down the outside of the thigh. As the problem progresses, the symptoms produce a limp when walking and stiffness in the hip joint. Eventually, the pain will also be present at rest and may even cause a problem with sleeping.



Diagnosis


How do doctors identify the problem?

The diagnosis of trochanteric bursitis begins with a history and physical examination. This is usually all that is necessary to make the

diagnosis. Your doctor will want to know when the pain began and which motions cause the pain. A physical examination will be done to determine how much stiffness you have in the hip and if you have a limp. Once this is done, X-rays will most likely be ordered to make sure that there are no other abnormalities in the hip.

X-rays will usually not show trochanteric bursitis. If X-rays are suggested they are to rule out other problems that may be causing your hip pain. Sometimes it is difficult to tell whether the pain you are suffering is from trochanteric bursitis or underlying arthritis of the hip joint. An X-ray may give more information about the condition of the hip joint itself.

An injection of a local anesthetic into the bursa can help your doctor diagnose trochanteric bursitis. If the injection removes the pain immediately, then the diagnosis is probably trochanteric bursitis. Most physicians will also add a bit of cortisone medication to the novocaine to help treat the condition at the same time.



Treatment

What can be done for the condition?

The treatment of trochanteric bursitis usually begins with simple measures. Treatment

becomes more involved if simple measures fail. The vast majority of patients with trochanteric bursitis will never require surgery.

Nonsurgical Treatment

Trochanteric bursitis is often treated successfully without surgery. Younger patients who have this condition because of overuse can usually be treated by reducing their activities or changing the way they do their activities. Combining this with an exercise program of stretching and strengthening and perhaps a brief course of anti-inflammatory medications will usually resolve the problem.

Your doctor may also prescribe sessions with a physical therapist. Treatments are used to calm inflammation and may include heat or ice applications. Therapists use hands-on treatment and stretching to help restore full hip range of motion. Improving strength and coordination in the buttock and hip muscles also enables the femur to move in the socket smoothly and can help reduce friction on the bursa. You may need therapy treatments for four to six weeks before full motion and function return.

If rehabilitation fails to reduce your symptoms, an injection of *cortisone* into the bursa may ease your symptoms and give temporary relief of the condition. Cortisone is a powerful anti-inflammatory medication. It can reduce swelling and pain when injected directly into the bursa. The injection will probably not cure the problem. But it may control the symptoms for months.

Surgery

Surgery is rarely needed to treat trochanteric bursitis. When all else fails and the pain is disabling, your doctor may recommend surgery. Several types of surgical procedures are available to treat trochanteric bursitis.

The primary goal of all procedures designed to treat this condition is to remove the thickened bursa, to remove any bone spurs that may have formed on the greater trochanter, and to relax

the large tendon of the gluteus maximus. Some surgeons prefer to simply lengthen the tendon a bit, and some prefer to remove a section of the tendon that rubs directly on the greater trochanter. Both procedures give good results.

Rehabilitation

What should I expect following treatment?

Nonsurgical Rehabilitation

Even if you don't need surgery, you may need to follow a program of rehabilitation exercises. Your doctor may recommend that you work with a physical therapist. Your therapist can create a program of stretching and strengthening for your hip. It is very important to maintain a balance of flexibility and strength

of the hip. You will probably progress to a home program within four to six weeks.

After Surgery

If you have surgery, your hip will be bandaged with a well-padded dressing. Physical therapy sessions may be needed for up to two months after surgery. The first few treatment sessions will focus on controlling the pain and swelling after surgery. You will then begin exercises that gradually stretch and strengthen the muscles around the hip joint. Your therapist will help you retrain these muscles to keep the ball of the femur moving smoothly in the socket. Your therapist will give you tips on ways to do your activities without straining the hip joint.

Notes

More about lateral hip pain

This booklet refers mainly to trochanteric bursitis but the term is loosely applied to lateral hip pain conditions. There are a number of possible cause of lateral hip pain and sorting out the diagnosis often requires more than one type of scan. The list of causes of lateral hip pain is long and though bursitis is common several other conditions frequently mimic the symptoms. Other common problems include tendon problems, hip joint pain, and lumbar (back) problems. Ultrasound alone may be insufficient to make a definite diagnosis but has the advantage of being useful to help guide targeted therapeutic injections. Ultrasound guided injections of corticosteroid and other substances directly into or around various structures can provide excellent symptomatic relief.

About Injections

Corticosteroid (cortisone) injections are part of managing the pain for this condition but do not change the long term outcome which is generally good. When the condition persists or there are significant underlying tendon problems then other substances such as POLIDOCANOL or rarely autologous blood (blood is taken from an arm vein and injected around a tear in a tendon) is sometimes recommended. This is done similar to corticosteroid injections but directly alters the healing response by causing sclerosis of small vessels and nerves (polidocanol) or stimulating healing (autologous blood). Whether this is appropriate depends on a host of factors including most importantly the ultrasound appearance in relation to the duration and type of symptoms.

For more about Injections for this condition see also : <http://www.driainduncan.com.au/tendinopathy/>